

Proxor®

100/6 & 200/6

(beclometasone/ formoterol)

Your health professional has prescribed Proxor® pMDI (pressurised metered dose inhaler) for you.

Proxor® provides relief from symptoms such as shortness of breath, wheezing and cough in people with asthma or chronic obstructive pulmonary disease (COPD). It also helps prevent asthma symptoms.

Proxor® contains two active substances: beclometasone dipropionate and formoterol fumarate dihydrate. They are inhaled through your mouth and delivered directly into your lungs. Together they make breathing easier.

Beclometasone dipropionate has an anti-inflammatory effect which helps reduce irritation in your lungs.

Formoterol fumarate dihydrate is a long acting bronchodilator which relaxes the muscles in your airways.



1. Cap. 2. Mouthpiece. 3. Base. 4. Push button. 5. Canister. 6. Dose counter





This guide is for patients who have been prescribed Proxor. It is designed to support you with your treatment and is not a substitute for the Package Leaflet that came with your medication.

Preparing your inhaler

Before first use, store your inhaler in a refrigerator (at 2-8°C) for a maximum of 18 months. Once open, use your inhaler for a maximum of three months and do not store above 25 °C. Before using the inhaler for the first time (or if you haven't used it for 14 days or more), hold your inhaler upright with the mouthpiece at the bottom and firmly depress the canister to release one puff.

Using your inhaler

Try to stand or sit in an upright position.



Don't forget

After use, replace the protective cap on your inhaler and check the dose indicator on the back. This shows how many puffs are left in your inhaler. When the dose indicator reaches 20, it's time to request another inhaler prescription.

If you find it difficult to use your inhaler while starting to breathe in, you may use the AeroChamber Plus spacer device. Ask your doctor, pharmacist or nurse about using a spacer.

The recommended dose of this medicine is one or two puffs twice daily. The maximum daily dose is 4 puffs.

To lower the risk of a fungal infection in your mouth or throat, rinse your mouth, gargle with water or brush your teeth after using your inhaler.

You should clean your inhaler once a week. For more information on this, visit proxor.co.uk.

Always use Proxor® exactly as your doctor, nurse or pharmacist has told you.

STADA
Thornton & Ross
Pharmaceuticals

Proxor®
100/6 & 200/6
(beclometasone/ formoterol)

Reporting of side effects. If you experience any side effects, talk to your doctor, nurse or pharmacist. This includes any possible side effects not listed in the package leaflet. You can also report side effects directly via the Yellow Card Scheme at yellowcard.mhra.gov.uk or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects, you can help provide more information on the safety of this medicine.